

Escalated Suicidal Rate of the Undergraduate Students of Bangladesh - Social Impact and Recommendations to Overcome

Mohammad Azizur Rahman^{1*}, Md. Touhidul Islam², Nabidur Rahman¹, Esrat Sultana³, Md. Sohel Parvez³, Shaikh Mirja Nurunnabi²

¹Department of Biochemistry and Molecular Biology, Jahangirnagar University, Bangladesh

²Department of Biochemistry and Molecular Biology, Shahjalal University of Science and Technology, Bangladesh

³Department of Microbiology, Jessore University of Science and Technology, Bangladesh

⁴Department of Microbiology, The Acme Laboratories Limited, Bangladesh

***Correspondence:** Mohammad Azizur Rahman, Department of Biochemistry and Molecular Biology, Jahangirnagar University, Bangladesh. E-mail: azizbmb@juniv.edu

Received: July 01, 2021; Accepted: July 16, 2021; Published: July 23, 2021

Copyright: © 2021 Rahman MA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Suicide has become one of the leading causes of death all over the world. It is considered a global public health problem and deviant behavior. People of different ages are committing suicide every day around the world for different reasons. University students, who are supposed to be the future of a country, are also accepting this path of death. Bangladeshi undergraduate students are also becoming the victims of suicide every day. In recent times, suicide ideation among university undergraduate students is increasing. This article reveals the prevalence of suicide among university undergraduate students and the factors associated with suicide ideation among university undergraduate students of Bangladesh. It also focuses on the impact of these suicides on the family and society. To overcome this grave situation, some precautionary steps have been recommended at the end of this article.

Key words: Affected behavior; Behavioral health; Mental disorder; Psychology; Public health; Suicide.

INTRUDUCTION

Suicide is the ninth leading cause of death in America leading to approximately 1.4% of the total deaths [1]. Several studies over the world have observed that university students face multiple problems ranging from psychological to academic and financial to career and social concerns [2]. It is also observed that these problems become harder with time and cannot be easily resolved to result in the need to seek some professional help. However, in developing countries like Pakistan, India, and Bangladesh, this scenario is more often due to a lack of social awareness. Accordingly, around 6 to 8% of students suffer through emotional issues during their university studies [3]. Though the exact number of students who suffer from emotional problems cannot be easily established, especially in developing countries. Researches have provided evidence that these students are at greater risk for academic and social challenges in a university environment and later in their practical adult

life [4]. Mental and behavioral disorders are estimated to account for 12% of the global burden of any disease. More than 40% of countries have no mental health policy and over 30% have no mental health program [5]. A number of factors are associated with suicidal ideation which includes depression, hopelessness, bullying, family divorce, parental psychopathology, parental death, low family income, interpersonal violence, psychiatric disorders, problematic technology use like smartphone addiction, Facebook or internet addiction, etc.[6-9]. In Bangladesh, the rate of suicide among university students is increasing day by day. It has become a serious issue that is needed to be addressed. But there is a lack of awareness regarding this suicide ideation among university students. Some proper steps may decline this rate significantly.

A study was conducted among 665 students of a university in Bangladesh by a classroom-based convenience sampling method. The study of the result

showed that the prevalence of suicidal ideation was 2.3% in the past 24 h, 4.8% in the past 15 days, 6.9% in the past month, 14.7% in the past year, and 61.1% in the lifetime. The results also stated that being separated from a partner, Facebook addiction, depression, anxiety, and stress were the risk factors of past-year Suicidal ideation [10]. In a retrospective study from the Bangladeshi press, a report found that most of the suicide victims are from public university students (30.4%). It found that 17 students were from a public university in a total of 56 Bangladeshi students suicide cases reported from January 2018 to June 2019 in the Bangladeshi print media [11]. A study in Jahangirnagar University among 407 university students aged between 18 and 27 years stated that 13.8% of the students have had suicidal ideation during the past 12 months. Several reasons were found to be responsible for this scenario. The prevalence of suicidal ideation was found more in the female than the male in the study [12]. Several studies show that female suicidal ideation is more in percentage than male. A study of suicidal ideation among rural elderly in Shandong, China shows that suicidal ideation varies gender-wise. It is in fact found more in female than the male. The reason of finding more suicidal ideation in female was stated because of having lower life satisfaction and negative life events which are more correlated with female than the male [13]. Other reasons that were found behind the suicidal ideation in the study conducted in Jahangirnagar University were being a fifth-year student, having a lower socioeconomic status, exposure to traumatic events, family suicide history, and depression. The lower socioeconomic status of individuals is more often influencing the university students to suicidal attempts. Most of the university students of Bangladesh belong to a lower or middle-class family. They often struggle to manage their education costs. Besides they must look after their families as well at a very early age. It often puts them under a huge amount of pressure. A cross-sectional study from a rural community of Bangladesh stated that the parental lower socio-economic position enhances the suicidal ideation rate among adolescents. Low monthly income and expenditure of the parents are associated with suicidal ideation of the adolescents and those who don't live with their parents are also found to have suicidal ideation [14]. A cross-sectional study from the Korean Community Health Survey 2013 confirmed that the suicide rate is higher among the people associated with lower socioeconomic status

[15]. On the other hand, Parental high expectations often put university students under tremendous pressure. Parents always seem to expect more from their children. They want their children to have great academic results. Besides, they often put their children under pressure through expecting their bright and luxurious future. The teachers also put a high load of academic pressure on them. Sometimes the university students do not find their teachers enough cooperative and helpful to share anything. These high expectations from their parents often put so much stress and anxiety on them. That is how the depression takes place in them and results in suicidal ideation. However, the impact of these suicides on family and society is very shocking. It enhances the breakdown of the family causing mental conflicts and depression among the family members. It creates traumatic stress that induces other family members to commit suicide. Suicide is a social disorder and an alarming situation for society. It creates anarchy in society and decreases social status. That is why it is considered as a defaming of society [16].

Some of the suicide prevention steps include raising awareness among university students regarding suicide, prevention programs are needed to arrange besides raising awareness, prevention programs should include student support centers, gatekeeper training towards mental health problems and suicide, screening, peer leadership training[17-19]. Besides, curriculum-based interventions and internet-based interventions may be beneficial in the prevention of university students' suicide [20]. finally, the university administration should look after the mental conditions of each student alongside their financial crisis.

CONCLUSION

Students' suicide is really a tremendous shock for both the family and society. This study will help to find out the exact causes of the suicide of University students. These study findings will help to address the relationship between suicide and student-teachers relationship and other factors. It is believed that the suicidal ideations of university students can be declined by taking meaningful steps. Some of them are also suggested in this study.

ACKNOWLEDGEMENT

Authors gratefully acknowledge the National Science and Technology (NST) fellowship 2019-20, provided to Md. Touhidul Islam by the Ministry of Science and Technology, Government of the People's Republic of Bangladesh.

REFERENCES

1. Beck AT, Brown GK, Steer RA, Dahlsgaard KK, Grisham JR. Suicide ideation at its worst point: a predictor of eventual suicide in psychiatric outpatients. *Suicide Life Threat Behav.* Spring 1999;29(1):1-9.
2. Rickwood D, Deane FP, Wilson CJ, Ciarrochi J. Young people's help seeking for mental health problems. *AeJAMH (Australian e-Journal for the Advancement of Mental Health).* 2005;4(3):218.
3. MacNeil Cheryl & Mead Shery. A Narrative Approach to Developing Standards for Trauma-Informed Peer Support. *Am J Eval.* 2005;26(2):231-244.
4. Eaton DK, Davis KS, Barrios L, Brener ND, Noonan RK. Associations of dating violence victimization with lifetime participation, co-occurrence, and early initiation of risk behaviors among U.S. high school students. *J Interpers Violence.* 2007;22(5):585-602.
5. Organization WHO and the World Health Report 2001-Mental Health: New Understanding, New Hope. World Health Organization. 2001.
6. Beautrais AL. Risk factors for suicide and attempted suicide among young people. *Aust N Z J Psychiatry.* 2000;34(3):420-436.
7. Blasco MJ, Vilagut G, Almenara J, Roca M, Piqueras JA, Gabilondo A, et al. Suicidal thoughts and behaviors: Prevalence and association with distal and proximal factors in Spanish university students. *Suicide Life Threat Behav.* 2019;49(3):881-898.
8. Franklin JC, Ribeiro JD, Fox KR, Bentley KH, Kleiman EM, Huang X, et al. Risk factors for suicidal thoughts and behaviors: A meta-analysis of 50 years of research. *Psychol Bull.* 2017;143(2):187-232.
9. Turecki G & Brent DA. Suicide and suicidal behaviour. *Lancet.* 2016;387(10024):1227-1239.
10. Mamun MA, Rayhan I, Akter K, Griffiths MD. Prevalence and Predisposing Factors of Suicidal Ideation Among the University Students in Bangladesh: A Single-Site Survey. *Int J Ment Health Addiction.* 2020.
11. Mamun MA, Siddique AB, Sikder MT, Griffiths MD. Student suicide risk and gender: A retrospective study from Bangladeshi press reports. *International Journal of Mental Health and Addiction.* 2020b.
12. Rahman ME, Saiful Islam M, Mamun MA, Moonajilin MS, Yi S. Prevalence and Factors Associated with Suicidal Ideation Among University Students in Bangladesh. *Arch Suicide Res.* 2020;18:1-10.
13. Lu L, Xu L, Luan X, Sun L, Li J, Qin W, et al. Gender difference in suicidal ideation and related factors among rural elderly: a cross-sectional study in Shandong, China. *Ann Gen Psychiatry.* 2020;19:2.
14. Begum A, Viitasara E, Soares J, Rahman AKMF, Macassa G. Parental socio-economic position and suicidal ideation among adolescents in Rural Bangladesh. *J Psychiatry Behav Sci.* 2018;4:1018.
15. Yeong Jun Ju, Eun-Cheol Park, Kyu-Tae Han, Jae Woo Choi, Jeong Lim Kim, Kyoung Hee Cho, et al. Low socioeconomic status and suicidal ideation among elderly individuals. *Int Psychogeriatr.* 2016;28(12):2055-2066.
16. Ara Mst, Uddin Md, Kabir Md. The causes of suicide and Impact of society in Bangladesh. *Int Res J Soc Sci.* 2016;5:25-35.
17. Arafat SMY, Mamun MAA. Repeated suicides in the University of Dhaka (November 2018): Strategies to identify risky individuals. *Asian J Psychiatr.* 2019;39:84-85.
18. Cusimano MD, Sameem M. The effectiveness of middle and high schoolbased suicide prevention programmes for adolescents: A systematic review. *Inj Prev.* 2011;17(1):43-9.
19. Masud MT, Mamun MA, Thapa K, Lee DH, Griffiths MD, Yang SH. Unobtrusive monitoring of behavior and movement patterns to detect clinical depression severity level via smartphone. *J Biomed Inform.* 2020;103:e103371.
20. Katz C, Bolton S, Katz LY, Isaak C, Tilston-Jones T, Sareen J, et al. A systematic review of school-based suicide prevention programs. *Depress Anxiety.* 2013;30(10):1030-45.